



Formosan Entomologist

Journal Homepage: entsocjournal.yabee.com.tw

New records of *Liriomyza* Mik (Agromyzidae: Diptera) leafminers from Indonesia **【Research report】**

印尼兩種潛蠅類之新紀錄 **【研究報告】**

M. B. MALIPATIL, P. M. RIDLAND A RAUF J. WATUNG, D. KANDOWANGKO
M. B. MALIPATIL, P. M. RIDLAND A RAUF J. WATUNG, D. KANDOWANGKO

*通訊作者E-mail: mallik.malipatil@dpi.vic.gov.au

Received: 2004/08/09 Accepted: 2004/09/27 Available online: 2004/12/01

Abstract

Two species of *Liriomyza* Mik (Agromyzidae), *L. katoi* Sasakawa and *L. yasumatsui* Sasakawa, collected from *Chrysanthemum* L. (Asteraceae) hosts, are newly recorded from Sulawesi, Indonesia. The taxonomic features that distinguish these 2 species from each other and from the major pest species, *L. trifolii* (Burgess), are discussed.

摘要

兩種斑潛蠅 *Liriomyza katoi* 斑潛蠅和 *L. yasumatsui* 斑潛蠅被從菊花寄主 (菊科 Asteraceae) 中採獲。這是印度尼西亞 Sulawesi 地區的全新記錄。兩種斑潛蠅的檢測以及與主要害蟲非洲菊斑潛蠅 *L. trifolii* 的外部形態的相似之處，都進行了討論並作了文獻記載。

Key words: *Liriomyza katoi*、*Liriomyza trifolii*、*Liriomyza yasumatsui*、*Chrysanthemum*、Asteraceae

關鍵詞: *Liriomyza katoi*、*Liriomyza trifolii*、*Liriomyza yasumatsui*、*Chrysanthemum*、Asteraceae

Full Text: [PDF\(0.27 MB\)](#)

下載其它卷期全文 Browse all articles in archive: <http://entsocjournal.yabee.com.tw>

New records of *Liriomyza* Mik (Agromyzidae: Diptera) leafminers from Indonesia

M. B. MALIPATIL, P. M. RIDLAND Department of Primary Industries Knoxfield, Private Bag 15, Ferntree Gully Delivery Centre, Vic. 3156, Australia

A RAUF Department of Plant Pests and Diseases, Faculty of Agriculture, Bogor Agricultural University, Bogor 16680, Indonesia

J. WATUNG, D. KANDOWANGKO Department of Plant Pests and Diseases, Faculty of Agriculture, Sam Ratulangi University, Manado, 95115, Indonesia

ABSTRACT

Two species of *Liriomyza* Mik (Agromyzidae), *L. katoi* Sasakawa and *L. yasumatsui* Sasakawa, collected from *Chrysanthemum* L. (**Asteraceae**) hosts, are newly recorded from Sulawesi, Indonesia. The taxonomic features that distinguish these 2 species from each other and from the major pest species, *L. trifolii* (Burgess), are discussed.

Key words: *Liriomyza katoi*, *Liriomyza trifolii*, *Liriomyza yasumatsui*, *Chrysanthemum*, Asteraceae

Introduction

The large agromyzid genus, *Liriomyza* Mik, with over 300 species worldwide (Spencer, 1973), is poorly represented in the Oriental region (Sasakawa, 1972). Fifteen species have been recorded from Taiwan (Shiao & Wu, 1995; Shiao & Wu, 2000) and at least twenty-five species from Japan (Sasakawa, 1992; 1993a, b, 1994; Iwasaki, 1993; Iwasaki *et al.*, 2000). Several of the species, including *L. huidobrensis* (Blanchard), *L. trifolii* (Burgess) and *L. sativae* Blanchard, are among the most destructive pests of horticultural crops, particularly vegetables and ornamentals (e.g. Spencer, 1973).

Liriomyza species currently known to occur in Indonesia include *L. brassicae* (Riley), *L. caulophaga* (Kleinschmidt), *L. chinensis* (Kato), *L. compositella* Spencer, *L. huidobrensis* and *L. sativae* (A. Rauf unpublished data; Shepard *et al.*, 1998; Spencer, 1961; Spencer, 1989). The pea leafminer, *L. huidobrensis*, has become a serious pest in highland vegetables in Java, Sumatra and South Sulawesi, causing yield losses as high as 60-70%. Another alien species, vegetable leafminer, *L. sativae*, has contributed to problems in lowland areas of Indonesia (Rauf *et al.*, 2000).

As part of an Australian Centre for International Agricultural Research

*Correspondence address
e-mail: mallik.malipatil@dpi.vic.gov.au

funded project on management and control of *Liriomyza* leafminers in Indonesia and Australia, a review of major pest species of leafminers present in different provinces and on different host plants in Indonesia is being conducted. A major aim of the review was to assess whether or not *L. trifolii* occurred in Indonesia. Leafminers collected on *Chrysanthemum* L. (Asteraceae) hosts at various localities in North Sulawesi, suspected to be *L. trifolii* based on major external morphological characters, were examined with emphasis on male genitalia which provide the most reliable diagnostic characters (Spencer, 1973). They were found not to belong to *L. trifolii* but to *L. katoi* Sasakawa and *L. yasumatsui* Sasakawa, both minor pest species hitherto unknown from Indonesia. A detailed redescription of these species is found in Shiao *et al.* (1991). Until now at least seven species of *Liriomyza* had been recorded from *Chrysanthemum* hosts, viz. *L. analis* (Rondani) from Italy, *L. spencerella* Valladares from Argentina, the widely distributed *L. huidobrensis*, *L. strigata* (Meigen) and *L. trifolii* (Spencer, 1990), *L. katoi* from Taiwan (Shiao *et al.*, 1991) and *L. ptarmicae* de Meijere from Europe, Canada, U.S.A. and Japan (Sasakawa, 1994).

All specimens examined in this study are deposited in the Victorian Agricultural Insect Collection, Department of Primary Industries, Knoxfield, Australia.

Taxonomy

Liriomyza katoi Sasakawa

Liriomyza katoi Sasakawa, 1961: Pac. Ins. 3: 400.

Diagnosis:

Face yellow; occiput black, black contiguous to ocellar triangle and eyes; antenna yellow with 3rd segment rounded; both vertical bristles on yellow background. Mesonotum shiny black; scutellum

yellow with both angles slightly brown-tinged; legs with coxae and femora yellow, tibiae and tarsi brown; wing length 1.5 mm in male, 1.6 mm in female, costa extending to M1+2. Abdominal tergites mostly shiny black. Male genitalia with phallus with characteristic large process on mesophallus and with rounded endophallus (Fig. 1), other details as in Shiao *et al.* (1991).

This species was until now recorded from Japan (on *Artemisia princeps*), Malaysia (on *Artemisia*), Taiwan (on *Chrysanthemum* sp.) (Shiao *et al.*, 1991; Sasakawa, 1994) and, in the present study, is newly recorded from Sulawesi (Indonesia) on *Chrysanthemum* sp.

Materials examined:

INDONESIA. North Sulawesi. 1 male, 3 females, Modoinding, 16-IX-2002, D.T. Sembel, on *Chrysanthemum*; 4 males, 2 females, Tomohon, 4-V-2003, D.T. Sembel, on *Chrysanthemum* sp.; 1 male, 3 females, no specific locality, 29-VI-2001, on *Chrysanthemum* sp.

Liriomyza yasumatsui Sasakawa

Liriomyza yasumatsui Sasakawa, 1972, Sci. Rep. Kyoto Univ. Agri. 24: 69.

Diagnosis:

As *L. katoi* except - occiput, ocellar triangle and arista brown, rest of head mainly yellow. Wing length 1.2 mm in male, 1.5 mm in female. Abdominal tergites brown. Male genitalia with phallus with characteristic large process on mesophallus, endophallus with sclerotised tubes narrowed (Fig. 2), other details as in Shiao *et al.* (1991).

This species was until now recorded from Taiwan (on *Artemisia* sp.) (Shiao *et al.*, 1991), Japan (on *Artemisia princeps*) (Sasakawa, 1994) and, in the present study, is newly recorded from Sulawesi (Indonesia) on *Chrysanthemum* sp.

Materials examined:




Fig. 1-3. Male genitalia of *Liriomyza* spp., lateral views; (1) *L. katoi*, (2) *L. yasumatsui*, (3) *L. trifolii*. Arrows indicate distiphallus. All figures to same scale. Scale line = 0.1 mm.

INDONESIA. North Sulawesi. 3 males, Modinding 16-IX-2002, D.T. Sembel, on *Chrysanthemum* sp.

Notes

Both these species may be readily confused with *L. trifolii* in characters such as yellow frons, yellow background under inner and outer vertical bristles, yellow hind margin of eye, yellow and rounded 3rd antennal segment, yellow scutellum and femora, and apex of costa extending to vein M1+2. However the latter differs from these species in having matt black mesonotum (shiny black in *L. katoi* and *L. yasumatsui*), and a characteristic male genitalia, particularly the structures of the phallus (Fig. 3).

Discussion

Correct identification of pest agromyzids has important implications for quarantine programs. Because male genitalia provide the most reliable taxonomic characters, identification of leafminers based on external morphological characters alone can lead to false identification of *L. trifolii*. When no adult males are available, it is valuable to complement morphological diagnoses with biochemical and molecular diagnoses, such as those developed by Central Science Laboratory in the United Kingdom (Central Science Laboratory, UK 2004) and in Japan (Miura *et al.*, 2004).

It is likely that there are more species of *Liriomyza* and other agromyzids waiting to be discovered in Indonesia, particularly in a poorly collected region such as North Sulawesi. Therefore there is a need for more detailed survey of agromyzid fauna of Indonesia on *Chrysanthemum* spp. (Asteraceae) hosts.

Acknowledgements

We thank Dr M. Sasakawa, 7-6-7 Korigaoka, Hirakata City, Osaka Pref., 573, Japan, for confirming the identifications, and Ms N. Endersby, DPI, Knoxfield, for critically reading the manuscript. The research was supported by ACIAR Project No. CS2/2000/090.

References

- Central Science Laboratory, UK.** 2004. Protocol for the diagnosis of quarantine organisms: *Liriomyza* spp. (*L. bryoniae*, *L. huidobrensis*, *L. sativae*, *L. trifolii*). www.csl.gov.uk/science/organ/ph/diagpro/liriomyza.pdf
- Iwasaki, A.** 1993. A new seed-feeding species of *Liriomyza* (Diptera, Agromyzidae) from Japan. *Japanese J. Entomol.* 61: 303-306.
- Iwasaki, A., K. Kasugai, R. Iwaizumi, and M. Sasakawa.** 2000. A newly recorded pest, *Liriomyza sativae* Blanchard in Japan. *Shokubutsu-boeki* 54: 12-17 (in Japanese).
- Miura, K., Y. Tagami, M. Ohtaishi, and A. Iwasaki.** 2004. Application of molecular techniques to distinguish *Liriomyza trifolii* from *L. sativae* (Diptera: Agromyzidae) on Tomato cultivation in Japan. *J. Econ. Entomol.* 97: 964-969.
- Rauf, A., B. M. Shepard, and M. W. Johnson.** 2000. Leafminers in vegetables, ornamental plants and weeds in Indonesia: surveys of host crops, species composition and parasitoids. *Int. J. Pest Management* 45: 1-10.
- Sasakawa, M.** 1961. A study of the Japanese Agromyzidae (Diptera). *Pac. Insects* 3: 307-472.
- Sasakawa, M.** 1972. Formosan Agromyzidae (Diptera). *Sci. Rep. Kyoto Pref. Univ. Agr.* 24: 43-82.
- Sasakawa, M.** 1992. Unrecorded leafminers (Diptera, Agromyzidae) from Japan. *Japanese J. Entomol.* 60: 558.
- Sasakawa, M.** 1993a. Notes on the Japanese Agromyzidae (Diptera), 1.

- Japanese J. Entomol. 61: 149-155.
- Sasakawa, M.** 1993b. Notes on the Japanese Agromyzidae (Diptera), 2. Japanese J. Entomol. 61: 341-354.
- Sasakawa, M.** 1994. Notes on the Japanese Agromyzidae (Diptera), 3. *Liriomyza*-miners on *Artemisa* spp. (Asteraceae). Japanese J. Entomol. 62: 55-64.
- Shepard, B. M., Samsudin, and A. R. Braun.** 1998. Seasonal incidence of *Liriomyza huidobrensis* (Diptera: Agromyzidae) and its parasitoids on vegetables in Indonesia. Int. J. Pest Management. 44: 43-47.
- Shiao, S. F, F. J. Lin, and W. J. Wu.** 1991. Redescription of four *Liriomyza* species (Diptera: Agromyzidae) from Taiwan. Chinese J. Entomol. 11: 65-74.
- Shiao, S. F., and W. J. Wu.** 1995. A new *Liriomyza* species from Taiwan (Diptera: Agromyzidae). Pan-Pacific Entomol. 71: 161-168.
- Shiao, S. F., and W. J. Wu.** 2000. *Liriomyza huidobrensis* (Blanchard), a newly invaded insect of economic importance to Taiwan (Diptera: Agromyzidae). Plant Prot. Bull. 42: 249-254.
- Spencer, K. A.** 1961. A synopsis of the Oriental Agromyzidae (Diptera). Trans. R. Entomol. Soc. Lond. 113: 55-100.
- Spencer, K. A.** 1973. Agromyzidae (Diptera) of economic importance. Series Entomologica volume 9. Dr. W. Junk B.V. Publisher, The Hague. 418 pp.
- Spencer, K. A.** 1989. Agromyzidae. In: N.L. Evenhuis (ed.). Catalog of the Diptera of the Australasian and Oceanian Regions. Bishop Museum Special Publication 86: 538-547.
- Spencer, K. A.** 1990. Host specialization in the world Agromyzidae (Diptera). Kluwer Academic Publishers, The Netherlands. 444 pp.
- Received: Aug. 9, 2004**
Accepted: Sep. 27, 2004

印尼兩種潛蠅類之新紀錄

M. B. MALIPATIL, P. M. RIDLAND Department of Primary Industries Knoxfield, Private Bag 15, Ferntree Gully Delivery Centre, Vic. 3156, Australia

A RAUF Department of Plant Pests and Diseases, Faculty of Agriculture, Bogor Agricultural University, Bogor 16680, Indonesia

J. WATUNG, D. KANDOWANGKO Department of Plant Pests and Diseases, Faculty of Agriculture, Sam Ratulangi University, Manado, 95115, Indonesia

摘 要

兩種斑潛蠅 *Liriomyza katoi* 斑潛蠅 和 *L. yasumatsui* 斑潛蠅被從菊花寄主 (菊科 Asteraceae) 中採獲。這是印度尼西亞 Sulawesi 地區的全新記錄。兩種斑潛蠅的檢測以及與主要害蟲非洲菊斑潛蠅 *L. trifolii* 的外部形態的相似之處, 都進行了討論並作了文獻記載。

關鍵詞 : *Liriomyza katoi*, *Liriomyza trifolii*, *Liriomyza yasumatsui*,
Chrysanthemum, Asteraceae