



Formosan Entomologist

Journal Homepage: entsocjournal.yabee.com.tw

Urban Forest Insect Pests and Their Management in Malaysia 【Review article】

馬來西亞的都市林害蟲和管理【綜合論述】

Chow-Yang Lee
李昭揚

*通訊作者E-mail: [✉ chowyang@usm.my](mailto:chowyang@usm.my)

Received: 2014/01/29 Accepted: 2014/02/06 Available online: 2014/02/01

Abstract

Urban forestry is an increasingly important field in Malaysia due to the greening efforts made by the government since 1997. Despite this, little is known about the insect pests affecting urban trees in Malaysia. This paper describes the common urban forest insect pests of Malaysia, the damages they cause, and the control methods commonly used against them.

摘要

由於自1997年起馬來西亞政府致力於綠化，都市林領域愈來愈重要；但有關為害馬來西亞都市林的害蟲資料不多。本文記述馬來西亞都市林常見害蟲、造成的損害及常用的防治方法。

Key words: urban forest, insect pest, Malaysia

關鍵詞: 都市林、害蟲、馬來西亞。

Full Text: [PDF\(0.36 MB\)](#)

下載其它卷期全文 Browse all articles in archive: <http://entsocjournal.yabee.com.tw>

Urban Forest Insect Pests and Their Management in Malaysia

Chow-Yang Lee

Urban Entomology Laboratory, Vector Control Research Unit, School of Biological Sciences, Universiti Sains Malaysia, 11800 Penang, Malaysia

ABSTRACT

Urban forestry is an increasingly important field in Malaysia due to the greening efforts made by the government since 1997. Despite this, little is known about the insect pests affecting urban trees in Malaysia. This paper describes the common urban forest insect pests of Malaysia, the damages they cause, and the control methods commonly used against them.

Key words: urban forest, insect pest, Malaysia

Introduction

The greening of urban Malaysia has primarily been focused on beautification of the urban landscape by horticulturists, landscapers, nursery workers, town planners, and architects, with minimal inputs from foresters. Because of this, very rarely is the term urban forestry being used (Sreetheran *et al.*, 2011). The first records of urban tree planting in Malaysia were in 1778 in Malacca (Koenig, 1894) and in 1802 in Penang (Burkill, 1966), where the Malay Paduak, *Pterocarpus indicus*, was planted. Because of its beauty and the ease of propagation, it was a popular tree for urban planting in Malaysia and Singapore until the 1990s (Sanderson *et al.*, 1997; Philip, 1999). However, the popularity of this species declined after the trees became infected

with *Fusarium* vascular wilt (*Fusarium oxysporum*). Currently, other indigenous or exotic species in Malaysia are replacing this species in urban planting programs.

In 1997, a nationwide tree planting program with the objectives of making Malaysia a garden nation was initiated (Nordin, 1997). With the support of city dwellers, private sectors, and government agencies, more than 400,000 trees and 6 million shrubs were planted (Sreetheran *et al.*, 2011). Trees were also grown as screens to protect against traffic dust and noise and to provide a nice view for people living next to motorways (Salleh *et al.*, 1990). Although the target of increasing tree number was reached, little was done to nurture and maintain the trees after they were planted. Regular tree inspection was often not performed, and insect pest problems and diseases were often not

[NOTE] The content of this review article was originally presented on 2013 International Symposium on Forest Health Management.

*Corresponding email: chowyang@usm.my

Urban Forest Insect Pests and Their Management in Malaysia 207

Table 1. Important pests of urban forests in Malaysia

Types of pests	Order	Family	Species	
Trunks and roots	Blattodea	Rhinotermitidae	<i>Coptotermes gestroi</i> <i>Coptotermes curvignathus</i> <i>Macrotermes gilvus</i>	
		Coleoptera	Termitidae	<i>Xystrocera festiva</i>
			Scarabaeidae	<i>Oryctes rhinoceros</i>
Leaves	Lepidoptera	Cossidae	<i>Squamura disciplaga</i>	
	Orthoptera	Acrididae	<i>Valanga nigricornis</i>	
		Lepidoptera	Papilionidae	<i>Chilasa clytia clytia</i>
	Lepidoptera	Hesperiidae	<i>Hasora cromus cromus</i>	
		Lymantriidae	<i>Orgyia postica</i>	
		Psychidae	Psychidae	<i>Pteroma pendula</i> <i>Mahasena corbetti</i> <i>Metisa plana</i> <i>Pagodiella hekmeyeri</i>
			Saturniidae	<i>Attacus atlas</i>
			Pyralidae	<i>Paliga damastesalis</i> <i>Sylepta derogata</i> <i>Neolithocolletis pentadesma</i>
		Coleoptera	Gracillariidae	<i>Apogonia</i> sp.
			Scarabaeidae	<i>Hypomeces squamosus</i>
Curculionidae	<i>Trachys</i> sp.			
Buprestidae	<i>Brontispa longgisima</i>			
Chrysomelidae	<i>Gynakothrips</i> sp.			
Phlaeothripidae	<i>Mictis longicornis</i>			
Sap suckers	Thysanoptera	Coreidae	<i>Aleurodicus disperus</i>	
	Hemiptera	Aleyrodidae	<i>Rastrococcus</i> sp.	
		Pseudococcidae	<i>Megatrioza vitiensis</i>	
Gall makers	Hemiptera	Psyllidae	<i>Quadrastichus erythrinae</i>	
	Hymenoptera	Eulophidae	<i>Eriophyes doctersi</i>	
	Acarina	Eriophyidae		

noticeable until the infestation became serious.

In this paper, I describe the common insect pests of urban trees in Malaysia and the problems that they cause. In addition, the common control approaches used against these insect pests in Malaysia are described.

Pests of urban trees in Malaysia and the damage they cause

A total of 28 species of insect and one species of acarine, covering 24 families and 8 orders, are commonly occurring

pests of urban forests in Malaysia (Table 1). The damage they cause can be divided into damage to 1) trunks, branches, and roots, 2) leaves, and 3) shoots and 4) loss of aesthetics due to the presence of the insects (Sajap, 2009).

Pests attacking tree trunks, branches, and roots

Subterranean termites are the primary pests of urban forests in Malaysia (Lee, 2007). Two species of termites, namely *Coptotermes curvignathus* and *C. gestroi*, are highly destructive to urban trees

Table 2. Prevalence of termite infestations among different tree species in a university campus in Penang, Malaysia

Species	No. inspected	% infested
<i>Acacia mangium</i>	16	87.5
<i>Casuarina equisetifolia</i>	13	84.6
<i>Fagraea fragrans</i>	13	84.6
<i>Araucaria excelsa</i>	12	75.0
<i>Tabebuia</i> sp.	13	69.2
<i>Albizia saman</i>	76	61.8
<i>Pterocarpus indicus</i>	47	61.7
<i>Swietenia</i> sp.	32	60.0
<i>Delonix regia</i>	35	57.1
<i>Araucaria cunninghamii</i>	11	54.5
<i>Pongamia pinnata</i>	30	50.0

(Cowie *et al.*, 1989). Normally, the damage caused by termites can be detected by the presence of mud tubes on the trunks of trees that have been attacked. Termites usually feed from the base of the tree leading to the inner side of the trunk, ultimately hollowing the tree and causing it to be structurally weakened. Newly attacked trees normally have leaves that are turning yellow and dropping. In ornamental palm trees, termites normally attack the spear region of the palm, and once the spear is destroyed, the tree dies because the spear is the growing vegetative part of the palm (UNEP, 2000). Another species of subterranean termite that attacks urban trees is *Macrotermes gilvus*. Normally, the damage it causes is limited to the tree bark; very rarely it will attack the inner trunk of mature live trees. Among the tree species planted on a university campus in Malaysia, the prevalence of termite infestations was found to vary with tree species (Table 2). Thus, some trees species are more susceptible to termite attack than others (Lee, 2009).

Xystrocera festiva is a longhorn beetle regularly found attacking moluca (peacock plume) and acacia trees in urban areas of Malaysia. The brown colored adult insect is between 35 and 40 mm long. The initial attack of this insect pest can be detected

by the presence of spotted sticky sap on the trunk that will eventually darken. The holes bored by the beetle are filled with wood debris and insect frass. The branches attacked by these beetles break easily, and infested trees eventually die.

Squamura disciplaga, a lepidopteran, is a major pest of shade trees such as casuarina, purple millettia, and Madras thorn. The larva bores into the trunk and feeds on the tree bark. It subsequently constructs galleries for protection using its silk, wood debris, and frass. Pupation occurs in the hole, and the pupa can be partially seen hanging out from the bored trunk. Unhealthy trees often experience additional microbial infections due to the holes bored by the insects.

Another important insect pest is the rhinoceros beetle, *Oryctes rhinoceros*. It is a large black beetle that often attacks urban palm trees. These beetles attack the developing fronds of the palm, move into the trunk, and then feed from the inside. Damaged fronds show triangular cuts. The larvae do not damage the trunk, although they do grow in the dead decaying trunk.

Pests attacking leaves

The pests that attack tree leaves are predominantly lepidopterans and coleopterans. In general, they are leaf miners, leaf

feeders, leaf-borers, leaf-curlers, or gall makers.

Bag worms (*Pteroma pendula*, *Metisa plana*, *Mahasena corbetti*, and *Pagodiella hekmeyeri*) are serious pests of ornamental urban palm trees in Malaysia such as the Manila palm and Macarthur palm. The biology of *P. pendula* is relatively unique. The adult female is apterous and vermiform like and spends its entire adulthood in the cocoon. The grey-colored adult male moth has a non-functional mouthpart. Palm leaves attacked by *P. pendula* looked rusty and often have pupae hanging below the leaves. Infestation often leads to loss of aesthetics and the shade characteristic of the palm trees. The damage caused by *M. plana* resembles that caused by *P. pendula*. The larvae of *M. plana* pupate in cocoons that are ~15 mm in length and are stuck to the leaf at an angle. *M. plana* also attacks other important urban trees such as purple millettia, moluca, acacia, and penaga lilin (*Mesua ferrea*).

The Atlas moth, *Attacus atlas* (L.), is the world's largest moth, with a wing span of up to 250 mm. It often feeds on young leaves of urban trees until the trees become leafless. Its common host trees are African mahogany, *Cinnamomum* spp., other species of amenity trees, and fruit trees (e.g., rambutan, guava, star fruit, and soursop) found in urban areas. Its caterpillar can reach 150 mm in length. The frass of the larvae can be spotted under infested trees.

Small pyralid moths, such as *Paliga damastesalis* and *Sylepta derogata*, are also pests of urban trees. *P. damastesalis* larvae feed on the leaf tissue of teak, often causing the leaves to turn brown and leaving only the veins. *S. derogata* larvae curl the leaves of *Erythrina* sp. and feed from inside. *Neolithocolletis pentadesma* is a small gracillariid moth whose larvae mine the leaves of moluca and other urban trees, creating a spotted membrane underneath the leaves. Repeated attacks by this species can cause premature

defoliation and lead to the death of branches. *Chilasa clytia clytia* larvae feed freely on the young leaves of *Cinnamomum* trees, especially newly planted ones, and can eventually lead to the death of the trees. *Hasora cromus cromus* larvae feed on the leaves of Indian beech (*Pongamia pinnata*).

There are several beetle species that attack leaves of urban trees. Adult scarab beetles, *Apogonia* sp., feed on the leaves of purple millettia, Bengal almond, moluca, and Indian beech and create holes in the leaves. Its larvae live in the soil and feed on grass roots and other small plants. The adults stay in the soil during the day and feed on leaves at night. The weevil, *Hypomeces squamosus*, feeds freely on the leaves of moluca, acacia, *Cinnamomum* spp., and yellow jade orchid tree (*Magnolia champaca*). Just like *Apogonia*, its larvae live in the soil and feed on grass roots. Buprestid larvae of the leaf miner, *Trachys* sp., feed on leaf tissue of urban shade trees such as Indian beech, causing the leaves to look spotty brown. The adult beetle then feeds on the entire leaf, except the hard veins. The chrysomelid *Brontispa longissima* attacks palm trees (such as betel nut and imperial palm) by feeding on their leaf tissues. The fronds of the palm turn brown and gradually become dry.

The locust *Valanga nigricornis* also feeds on amenity trees and attacks plants in nurseries and newly planted trees. However, it is not a major insect pest of urban trees in Malaysia.

Sapsuckers

Sapsuckers are insects with piercing/sucking mouthparts. Four pest species from the order Thysanoptera and Hemiptera are commonly found attacking urban trees in Malaysia. Besides causing injury to the trees, these pests can also serve as vectors of plant viruses.

The thrip, *Gynakothrips* sp., which measures 0.5 mm long, often attacks

shade trees from the genus *Ficus*. Affected leaves are often twisted, spotted, and dark brown in color. The coreid bug, *Mictis longicornis*, sucks the sap from young shoots. When disturbed, it releases a pungent liquid that irritates the eyes. The whitefly, *Aleurodicus disperus*, feeds by tapping into the phloem of plants, introducing toxic saliva and decreasing the plants' overall turgor pressure. The mealybug, *Rastrococcus* sp., attacks many species of urban trees including *Plumeria*. The honeydew of the bug causes the leaves to turn dark and become covered by a layer of soot, which decreases the aesthetics of the trees.

Gall makers

Three species of pests in Malaysia are known to attack the meristematic plant tissues of urban trees causing unusual deformations known as galls. The leaf hopper, *Megatrioza vitiensis*, normally causes galls on *Eugenia papillose*, *Cinnamomum* spp., and Indian rose chestnut trees. Eggs are laid under the leaf, and the nymphs suck the leaf sap. This causes the upper surface of the leaf to form a gall. Erythrina gall wasp (*Quadrastichus erythrinae*) is an important gall-making insect on coral trees in Malaysia such as *Erythrina variegata*, *Erythrina crista-galli* and *Erythrina glauca*. The midge-like adult wasp will insert their eggs into young leaves and stem tissues. Wasp larvae hatch and develop in plant tissues inducing gall formation. Normally there is a wasp per gall cell, but could be more under heavy infestation. Larvae pupate within the gall, and the adults emerge by cutting exit holes through the galls. The other species is an acarine, *Eriophyes doctersi* that normally attacks the leaves of *Cinnamomum iners*.

Control of pests of urban forests in Malaysia

Chemical control remains the most common method used against pests of urban forests in Malaysia (Sajap, 2009). Table 3 provides a summary of the common control methods used. Trees infested with subterranean termites such as *C. gestroi* and *C. curvignathus* are normally drenched with insecticide (e.g., chlorpyrifos, fipronil, and imidacloprid). Before drenching, the mud tubes are removed from the trunk and a trench is dug along the base of the tree. Bait can also be used against these two species. For example, in one study in-ground termite monitoring stations were installed, and when the stations became infested, termite baits containing a chitin synthesis inhibitor (e.g., hexaflumuron, chlorfluazuron, and bistrifluron) were placed in the infested stations (Lee, 2001). For higher termites such as *M. gilvus*, mounds are located and excavated and the remains are drenched with a pyrethroid. For beetle pests attacking tree trunks, imidacloprid and azadirachtin can be injected into the trunks. However, trees that are seriously infested are destroyed to avoid further infestation of other healthy trees. For the rhinoceros beetle, the fungal agent *Metarhizium anisopliae* can be applied to the piles of infested trunks. To manage *S. disciplaga*, dimethoate or fenitrothion are injected into the hole bored by the insect pest.

When controlling lepidopteran pests attacking leaves, the systematic insecticide carbofuran and chemical sprays often are used. Systemic insecticides have proven to be effective against bagworms such as *P. pendula* and *M. plana*. Microbial agents such as *Bacillus thuringiensis* can also be used through application into the affected trees. Injection of tree branches with monocrotophos can also be used to manage bagworms. To manage coleopteran pests attacking leaves, leaves often are sprayed with pyrethroid and organophosphate sprays. Chlorpyrifos can be used to control insect larvae in the soil.

Table 3. Common control methods used against pests of urban forests in Malaysia

Species	Treatment option	
	Chemical/others	Biological
<i>Coptotermes gestroi</i>	Drenching, termite baiting	–
<i>Coptotermes curvignathus</i>	Drenching, termite baiting	–
<i>Macrotermes gilvus</i>	Excavation of mound and drenching	–
<i>Xystrocera festiva</i>	Chemical injection into the trunk	–
<i>Oryctes rhinoceros</i>	Pheromonal trap, chemical injection	<i>Metarhizium</i> spray
<i>Squamura disciplaga</i>	Chemical injection into the trunk	–
<i>Valanga nigricornis</i>	Malathion, methamidophos or monocrotophos sprays	–
<i>Chilasa clytia clytia</i>	Abamectin spray	Bt spray
<i>Hasora cromus cromus</i>		Bt spray
<i>Orgyia postica</i>		Bt spray
<i>Pteroma pendula</i>	Systemic insecticide carbofuran	Bt spray
<i>Mahasena corbetti</i>	Systemic insecticide carbofuran	Bt spray
<i>Metisa plana</i>	Systemic insecticide carbofuran	Bt spray
<i>Pagodiella hekmeyeri</i>	Acephate spray	–
<i>Attacus atlas</i>	Malathion or cypermethrin sprays	Bt spray
<i>Paliga damastesalis</i>		Bt spray
<i>Sylepta derogata</i>	Permethrin or carbaryl sprays	–
<i>Neolithocolletis pentadesma</i>	Injection of acephate	–
<i>Apogonia</i> sp.	Light trap, cypermethrin, or trichlorfon sprays on leaves, chlopyrifos on soil	–
<i>Hypomeces squamosus</i>	Acephate, carbaryl or methamidophos sprays	–
<i>Trachys</i> sp.	Pyrethroid sprays	–
<i>Brontispa longgisima</i>	Systemic insecticide such as carbofuran	–
<i>Gynakothrips</i> sp.	Removal of affected branches, and deltamethrin or fipronil sprays	–
<i>Mictis longicornis</i>	Acephate or dimethoate sprays	–
<i>Aleurodicus disperus</i>	Removal of affected leaves, white oil, profenofos or imidacloprid sprays.	–
<i>Rastrococcus</i> sp.	Removal of affected leaves, white oil or malathion sprays	–
<i>Megatrioza vitiensis</i>	Imidacloprid, malathion, permethrin, spinosad sprays	–
<i>Quadrastichus erythrinae</i>	Foliar application of dimethoate, imidacloprid, fipronil and cypermethrin.	–
<i>Eriophyes doctersi</i>	Imidacloprid, malathion, permethrin, spinosad sprays	–

To treat infestation with sapsuckers, affected branches and leaves are removed before the application of insecticides (e.g., deltamethrin, fipronil, acephate, dimethoate, profenofos, and imidacloprid). White oil can be used against mealybugs and

whiteflies. Lastly, foliar sprays (dimethoate, imidacloprid, malathion, permethrin, and spinosad) applied to affected leaves often are used against gall makers.

Acknowledgments

I would like to thank Ahmad Said Sajap (Universiti Putra Malaysia) and Arthur Y. C. Chung (Sabah Forest Department) for providing a lot of valuable information that has made this paper possible.

References

- Burkill IH.** 1966. A dictionary of the economic products of the Malay Peninsula. Volumes 1 and 2. Ministry of Agriculture and Cooperation, Kuala Lumpur, Malaysia.
- Cowie RH, Logan JWM, Wood TG.** 1989. Termite (Isoptera) damage and control in tropical forestry with special reference to Africa and Indo-Malaysia: a review. *Bull Entomol Res* 79: 173-184.
- Koenig JG.** 1894. Journal of a voyage from India to Siam and Malacca in 1779. *J Straits Branch Roy Asiatic Soc* 26: 58-201.
- Lee CY.** 2001. Subterranean termite pests and their control in the urban environment in Malaysia. *Sociobiology* 40: 3-9.
- Lee CY.** 2007. Perspective in urban insect pest management in Malaysia. Penang, Malaysia: Vector Control Research Unit, Universiti Sains Malaysia. 106 pp.
- Lee CY.** 2009. Subterranean termites of Universiti Sains Malaysia. pp 33-42. In: Ismail A, Mansor M, Yahya K, Baharuddin MH, Lee CY (eds). *The ecosystem, landscape and heritage of Universiti Sains Malaysia*. Universiti Sains Malaysia.
- Nordin AR.** 1997. Managing the garden city. pp 73-84. In: Othman MT, Mustafa Kamal MS, Noorizan M, Nordin AR, Abdul Aziz O (eds). *Ke Arah Negara Taman: Wawasan dan Cabaran*. Institute of Landscape Architect Malaysia, Shah Alam, Malaysia.
- Philip E.** 1999. Wilt disease of angkana (*Pterocarpus indicus*) in Peninsular Malaysia and its possible control. *J Trop Forest Sci* 11: 519-527.
- Sajap AS.** 2009. Serangga perosak pokok ameniti. Jabatan Landskap Negara, Kementerian Perumahan dan Kerajaan Tempatan (in Malay).
- Salleh MN, Wong YK, Ng FSP.** 1990. The tropical garden city – Its creation and maintenance. *Malayan Forest Record*, No. 33. Kuala Lumpur: Forest Research Institute of Malaysia. 19 pp.
- Sanderson FR, Fong YK, and Saiful A.** 1997. A *Fusarium* wilt (*Fusarium oxysporum*) of angkana (*Pterocarpus indicus*) in Singapore. *J Arboricul* 21: 205-214.
- Sreetheran M, Adnan M, Khairil Azuar AK.** 2011. Street tree inventory and tree risk assessment of selected major roads in Kuala Lumpur, Malaysia. *Arboricul Urban Forest* 37: 226-235.
- UNEP (United Nation Environment Programme).** 2000. Finding alternatives to persistent organic pollutants (POPs) for termite management. UNEP. <http://www.unep.org/hazardoussubstances/Portals/9/Pesticides/Alternatives-termite-fulldocument.pdf> (retrieved on 10 October 2013).

Received: January 29, 2014

Accepted: February 6, 2014

馬來西亞的都市林害蟲和管理

李昭揚

馬來西亞理科大學生物系都市昆蟲研究室

摘 要

由於自 1997 年起馬來西亞政府致力於綠化，都市林領域愈來愈重要；但有關為害馬來西亞都市林的害蟲資料不多。本文記述馬來西亞都市林常見害蟲、造成的損害及常用的防治方法。

關鍵詞：都市林、害蟲、馬來西亞。

[註] 本文內容原於「2013 年森林健康之管理與經營國際研討會」中發表。

*論文聯繫人

Corresponding email: chowyang@usm.my